

RESEARCH ARTICLE

Is atopic sensitization associated with indicators of early vascular ageing in adolescents?

Karsten Königstein^{1*}, Denis Infanger¹, Randi Jacobsen Bertelsen^{2,3}, Ane Johannessen^{3,4}, Ulrike Waje-Andreassen⁵, Arno Schmidt-Trucksäss¹, Cecilie Svanes^{3,4‡}, Julia Dratva^{6,7‡}

1 Department of Sport, Exercise and Health, Division Sports and Exercise Medicine, University of Basel, Basel, Switzerland, **2** Department of Clinical Science, University of Bergen, Bergen, Norway, **3** Department of Occupational Medicine, Haukeland University Hospital, Bergen, Norway, **4** Centre for International Health, Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway, **5** Department of Neurology, Haukeland University Hospital, Bergen, Norway, **6** Medical Faculty, University of Basel, Basel, Switzerland, **7** ZHAW, School of Health Professions, Winterthur, Switzerland

‡ These authors are last authors on this work.

* k.koenigstein@unibas.ch



OPEN ACCESS

Citation: Königstein K, Infanger D, Jacobsen Bertelsen R, Johannessen A, Waje-Andreassen U, Schmidt-Trucksäss A, et al. (2019) Is atopic sensitization associated with indicators of early vascular ageing in adolescents? PLoS ONE 14(8): e0220198. <https://doi.org/10.1371/journal.pone.0220198>

Editor: Rudolf Kirchmair, Medical University Innsbruck, AUSTRIA

Received: March 25, 2019

Accepted: July 10, 2019

Published: August 15, 2019

Copyright: © 2019 Königstein et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The study population is from a restricted geographical area, and in order to ensure full anonymity only the main variables of the final analyses are provided. We confirm that the data file provided constitutes the minimal dataset necessary to replicate the findings of your study in their entirety. However, we are open to making the full parameters available for scientific analysis on request, provided that the respective research institution proves handling of the data strictly in accordance with ethical regulations (written ethics

Abstract

Background

Chronic systemic inflammation accelerates early vascular ageing. Atopic sensitization and allergic diseases may involve increased inflammatory activity. This study aimed to assess whether atopic sensitization and allergic diseases were associated with altered vascular biomarkers in Norwegian adolescents.

Methods

Distensibility coefficient of the common carotid arteries, carotid intima-media thickness and atopic sensitization (serum total and specific IgEs) were assessed in 95 Norwegian adolescents, who participated in the RHINESSA generation study. Symptoms of allergic disease were assessed by an interviewer-led questionnaire.

Results

Atopic sensitization was found in 33 (34.7%) of the adolescents. Symptomatic allergic disease was found in 11 (33.3%) of those with atopic sensitization. Distensibility coefficient of the common carotid arteries appeared to be lower in participants with atopic sensitization than in those without ($46.99 \pm 8.07 \times 10^{-3} / \text{kPa}$ versus $51.50 \pm 11.46 \times 10^{-3} / \text{kPa}$; $p > 0.05$), while carotid intima-media thickness did not differ between these groups ($0.50 \pm 0.04 \text{ mm}$ versus $0.50 \pm 0.04 \text{ mm}$; $p > 0.05$). Crude, as well as age- and sex-adjusted multiple regression, revealed no significant association, neither of atopic sensitization nor of allergic disease, with distensibility coefficient of the common carotid arteries and carotid intima-media thickness.

protocol, full compliance with the Declaration of Helsinki). Data requests can be made to Department of Occupational Medicine, Haukeland University Hospital, 5021 Bergen (datamanager in charge: Rebnord, Eivind Aksnes <eivind.aksnes.rebnord@helse-bergen.no>).

Funding: Overview of funding sources for the RHINESSA study can be found on: <https://helse-bergen.no/seksjon/RHINESSA/Documents/Funding%20RHINESSA.pdf>. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

Our results do not support the assumption of an adverse impact of atopic sensitization and/or allergic disease on distensibility coefficient of the common carotid arteries and carotid intima-media thickness in Norwegian adolescents. Further research is necessary to study whether the clinical severity of allergic diseases might be more important than the status of allergic disease or atopic sensitization.

Introduction

Evidence is growing, that inflammatory processes play an important role in atherogenesis, promoting the risk of cardiovascular diseases [1]. Possible pathophysiological links between inflammation and vascular damage were previously described [2–8]. One of the most investigated mechanisms is the oxidative modification of LDL, which leads to foam cell formation and development of lesions in the vascular wall [2, 3]. Wang et al. observed a stimulated arterial cell apoptosis and cytokine expression in humans and mice by elevated serum IgE levels [4]. This might be preceded by decreased serum-levels of low-affinity IgE receptor-positive B cells, as observed after coronary artery bypass graft surgery [5]. A possible link between chronic inflammatory activity caused by atopic sensitization and atherosclerosis might be an elevated activity of mast cells, which leads to multiple effects in the vascular wall, promoting development and vulnerability of atherosclerotic lesions [6]. The influence of childhood exposure to several pro-inflammatory risk factors on vascular health in adult life has previously been shown, as well as the relevance of childhood exposure to cardiovascular risk factors for the later development of atherosclerosis [9–13]. Repeated bacterial or viral infections, obesity and diabetes mellitus are strong promoters of increased carotid intima-media thickness (cIMT) in children by means of chronically elevated inflammatory activity [14–16]. So far, a possible association of chronic systemic inflammation related to atopic sensitization with cIMT and other biomarkers of early vascular ageing has been analyzed mainly in adult populations [17, 18]. However, a few studies suggested that allergic diseases might contribute to early vascular ageing already in early childhood [19, 20].

Atopic sensitization, independent of its clinical penetrance, involves chronic systemic hyperinflammation. Hence, we hypothesized that atopic sensitization might contribute to early vascular ageing in young people. The primary aim of this study was to investigate a possible association of atopic sensitization, independent of its clinical significance, with the distensibility coefficient (DC) of the common carotid arteries and cIMT, which are indicators of early vascular ageing. Our second aim was to investigate whether the clinical manifestation of atopic sensitization might be associated with these parameters. Hence, we also analyzed the association of allergic disease with DC and cIMT.

Methods

Study population

All Norwegian offspring aged 10 to 18 years of ECRHS Bergen participants were invited to participate in the prospective RHINESSA generation study (Respiratory Health In Northern Europe, Spain and Australia, see www.rhinessa.net, Fig 1). Of the 285 offspring 125 had parental consent for clinical investigation and were screened for eligibility. Exclusion criteria were a recent operation, an acute infection, diabetes mellitus or other chronic inflammatory diseases




Fig 1. Participant recruitment process. * n = 2 with type 1 diabetes mellitus. Further predefined exclusion criteria were recent operation, an acute infection, diabetes mellitus type 2, any chronic inflammatory diseases unrelated to atopy, severe heart disease and pregnancy; & No participant's consent for blood tests.

<https://doi.org/10.1371/journal.pone.0220198.g001>

unrelated to atopy, severe heart disease or pregnancy. Overall, we excluded two candidates because of diabetes mellitus type 1. Of the remaining 123 participants, 21 had no test of immunological total or specific IgE, because they had not agreed for blood analysis. Furthermore, seven participants' ultrasound images did not meet the predefined quality criteria (see [S1](#) and [S2 Files](#) for further details). Therefore, 95 participants were available for main analysis ([Fig 1](#)).

Data analysis was performed in accordance with the Declaration of Helsinki and approved by the Regional committee for Medical and Health Research Ethics, Western Region (REC West 2012/1077). Written informed consent was retrieved prior to participation (from parents if the offspring was below 16 years of age, from the offspring themselves if 16 years or older).

Questionnaires

Extensive information on respiratory health, allergic diseases, general health and environmental exposures were assessed by a web-based questionnaire, covering all possible covariates for the analysis: parents' atopy status, physical activity, frequent exposure to smoking (either active smoking or exposure to regular parental smoking at home), modality of birth (caesarean section versus natural birth) and preterm birth.

An interviewer-led questionnaire before clinical examination assessed respiratory symptoms during the last months and specifically during the last three days and current medication. Participants were asked to bring any regular or emergency medication to the study centers.

Clinical examinations

Based on interview data on hours since having smoked or consumed food and drinks, medication use, and current infections, no participants were excluded from specific examinations. Afterwards, we conducted spirometry, FeNO analysis, analysis of total and specific IgE and anthropometric measures (see [S1 File](#) for detailed information).

Main predictors: Atopic sensitization and allergic disease

Atopic sensitization was defined as a positive total or specific IgE towards inhalant allergens (house dust mite, cat, Timothy (grass, birch, and Cladosporium). Allergic disease was defined as atopic sensitization plus two of the following clinical criteria: allergic rhinitis, atopic eczema, food allergy, allergic bronchial asthma or frequent use of doctor-prescribed antihistaminic medication (see [S1 File](#) for further details).

Ultrasonographic examination and main outcomes: DC and cIMT

ECG derived heart rate as well as systolic and diastolic blood pressure were obtained simultaneously during the ultrasonographic assessment of DC and cIMT. Blood pressure was measured on the left upper arm with an OMRON 705 IT-IS Automatic-IS device just before the examination was started and after ten minutes of rest in a sitting position. The appropriate cuff size was determined by measuring the upper arm circumference.

The procedure of DC and cIMT measurement was performed by two trained field workers, using an ultrasound instrument (UF-870, Fukuda Denshi Co. Ltd., Tokyo, Japan) with a LA38 5–16 MHz linear probe. Temporal resolution was 10.47 ms per frame. Data assessment was conducted using an automated wall-detection software, as previously described and in accordance with current recommendations to ensure acceptable data quality (see [S2 File](#) for further details; [21, 22]). In the pre-study training examinations intraobserver variability was 7.2 and 7.9%, respectively, and, thus, similar to typical values [23]. Interobserver variability was 10.7% and, thus, slightly higher than previously reported values [23]. Near- and far-wall cIMT as well as end diastolic and peak systolic outer lumen diameter were obtained from four examination planes (bilateral common carotid artery horizontal plane and ear-to-ear plane, respectively) [21]. Afterwards, a mean DC [10^{-3} /kPa] and mean cIMT [mm] were calculated for each participant and used for further statistical analysis (see [S2 File](#) for further details). Validity,

reliability and clinical predictive value of DC and cIMT in children and adolescents have been shown previously [24–27].

Statistical analyses

Data analysis was performed using SPSS version 25.0 for Windows (SPSS Inc., Chicago, Illinois, USA) and R version 3.5.0 for Windows (R Foundation for Statistical Computing, Vienna, Austria). Descriptive analysis included means, standard deviations (SD), minimum and maximum values. The level of significance was set at $p \leq 0.05$; estimated effects were reported with 95% confidence intervals (95% CI).

Unadjusted (crude) linear regression models, as well as age- and sex-adjusted multiple regression models were applied to analyze associations of DC and cIMT with atopic sensitization and allergic diseases.

Complete data were available for 95 (81.1%) of 117 participants. In the remaining 22 cases parental or participants' consent for IgE analyses was not given. We used multiple imputation by chained equations using the “mice” package in R (version 3.0.3) to impute the missing data [28, 29]. Specifically, we imputed 300 datasets with 10 iterations each. Convergence and distribution of imputed values were assessed graphically. We applied predictive mean matching for imputation of continuous variables, logistic regression for binary variables and ordered logistic regression for ordered categorical variables [30]. The results from the regression models based on the imputed datasets were pooled using Barnard-Rubin adjusted degrees of freedom for small samples [31].

We repeated all statistical analyses with height-related standard deviation scores (SDS) of DC and cIMT, because height seems to be a strong determinant for vascular wall properties in childhood and adolescence [25]. Sensitivity analyses were performed including factors, which are more or less controversially discussed in literature as possibly being relevant for early vascular ageing and also the risk of atopic sensitization. These factors are current exposure to smoking (passive and active) [32, 33], physical activity [34], preterm birth [35] and delivery by caesarean section [36]. None of these additional analyses resulted in significantly different outcomes and, therefore, they are not presented.

Results

Basic characteristics of the study population

General empiric and vascular characteristic were comparable in participants with and without atopic sensitization (Table 1). Atopic sensitization was found in 33 (34.7%) of all participants with available blood samples. Of these participants, allergic disease was found in 11 (33.3%) individuals. Nine of them took antiallergic medication on a regular basis at the time of examination. Mean DC was $46.99 \pm 8.07 * 10^{-3}$ /kPa in the group with atopic sensitization and $51.50 \pm 11.46 * 10^{-3}$ /kPa in the group without atopic sensitization. Mean cIMT was 0.50 ± 0.04 mm in both groups.

Association of atopic sensitization with biomarkers of early vascular ageing

Neither crude comparison of DC and cIMT, nor the age- and sex-adjusted multivariate regression model, indicated a significant association of atopic sensitization with these parameters (Table 2). However, DC tended to be lower in participants with atopic sensitization than in those without.

Association of allergic disease with biomarkers of early vascular ageing

Neither crude comparison of DC and cIMT, nor the age- and sex-adjusted multivariate regression model, indicated a significant association of allergic disease with DC and cIMT (Table 2).

Table 1. Population characteristics.

	Mean (standard deviation; min/max) / n (%)		
	Atopic sensitization (n = 33; 34.7%)	No atopic sensitization (n = 62; 65.3%)	p-value [#]
<i>Empiric characteristics</i>			
Age [years]	15.3 (±1.7; 12.4/18.5)	15.1 (±2.7; 9.8/18.7)	ns
Female	16 (48.5%)	32 (51.6%)	-
Height [cm]	169 (±9.2; 148/189)	166 (±13.7; 130/188)	ns
BMI [kg/m ²]	21.4 (±3.3; 16.0/28.7)	21.0 (±4.1; 14.7/38.2)	ns
BMI-SDS ^{&} [37]	0.58 (±1.23; -1.24/3.27)	0.43 (±1.64; -2.48/6.97)	ns
<i>Vascular characteristics</i>			
DC [10 ⁻³ /kPa]	46.99 (±8.07; 28.79/63.97)	51.50 (±11.46; 29.30/95.83)	ns
DC-SDS [#] [25]	-0.77 (±0.62; -2.37/0.27)	-0.48 (±0.78; -2.51/2.26)	ns
cIMT [mm]	0.50 (±0.04; 0.42/0.59)	0.50 (±0.04; 0.42/0.62)	ns
cIMT-SDS [#] [25]	2.30 (±0.88; 0.40/4.19)	2.31 (±0.85; 0.69/4.53)	ns
BP _{sys} [mmHg]	108 (±8.1; 92/123)	107 (±8.4; 91/128)	ns
BP _{dia} [mmHg]	58 (±4.7; 49/73)	62 (±6.5; 52/90)	0.01
Heart rate [bpm]	68 (±19.2; 48/143)	69 (±18.8; 45/170)	ns

BMI = Body mass index; BMI-SDS = Standard deviation score of BMI; DC = Distensibility coefficient of the common carotid arteries; DC-SDS = Standard deviation score of DC; cIMT = Carotid intima-media thickness; cIMT-SDS = Standard deviation score of cIMT; BP_{sys} = Blood pressure, systolic; BP_{dia} = Blood pressure, diastolic.

[#]T-test derived differences between subpopulations with and without atopic sensitization were considered significant, if two-tailed p ≤ 0.05 (ns = p > 0.05).

[&]Norwegian reference population.

*Mainly European reference population.

<https://doi.org/10.1371/journal.pone.0220198.t001>

Discussion

Mean DC tended to be lower in participants with atopic sensitization than in those without. However, atopic sensitization revealed no significant association with DC and cIMT in this study population of Norwegian adolescents. Further, no significant associations of clinically apparent allergic diseases with DC and cIMT were identified.

In 2015, Evelein et al. found an increased cIMT in five-year old children with several clinical forms of allergies but no changes in arterial distensibility and elasticity [20]. They

Table 2. Multiple linear regression analysis of associations of atopic sensitization and allergic disease with DC and cIMT.

Model		β	lower 95% CI	upper 95% CI	p
Atopic sensitization ^{&}	DC crude	-3.10	-8.80	2.60	0.28
	DC adjusted [*]	-3.04	-7.99	1.90	0.22
	cIMT crude	0.004	-0.01	0.02	0.66
	cIMT adjusted [*]	0.004	-0.01	0.02	0.67
Allergic disease [#]	DC crude	-1.94	-10.42	6.53	0.64
	DC adjusted [*]	-1.01	-8.89	6.87	0.80
	cIMT crude	0.009	-0.02	0.04	0.54
	cIMT adjusted [*]	0.008	-0.02	0.04	0.60

[&]Atopic sensitization = any positive total or specific IgE in the serological analyses (dermatophagoides, cat, birch, timothy grass, cladosporium)

[#]Allergic disease = atopic sensitization plus two of the following clinical criteria: allergic rhinitis, atopic eczema, food allergy, allergic bronchial asthma or frequent use of doctor-prescribed antihistaminic medication

*Models adjusted for age and sex. DC = Distensibility coefficient of the common carotid arteries; cIMT = Carotid intima-media thickness; β = Estimated effect; CI = Confidence interval; level of significance set at p ≤ 0.05.

<https://doi.org/10.1371/journal.pone.0220198.t002>

concluded that allergies are associated with arterial changes in young children. However, our data do not support their cIMT findings. Possibly, age as well as timing and severity of clinical manifestations of atopy might play a role. Yet, the tendency towards a lower DC in our participants with atopic sensitization might be a very early sign for a chronic subclinical impact of atopic sensitization on vascular ageing. Helpful markers of the severity of systemic inflammatory activity (i. e. oxLDL, high-sensitivity C-reactive protein, soluble interleukin-2 receptor, eosinophil cationic protein) were not available in both studies. However, we analyzed total and specific IgE, which are valid markers for qualitative assessment of atopic sensitization but do not give information about the severity of systemic inflammatory activity [38]. We assume that the inflammatory activity in our study population might have been somewhat heterogeneous, which would explain the lack of association of atopic status and allergic disease with DC and cIMT. Whether the strength of such an association might depend on the severity of systemic inflammatory activity needs to be investigated in larger prospective studies including more information about current and cumulative lifetime systemic inflammatory activity.

Another study found an adverse influence of repeated episodes of common childhood infectious diseases on cIMT [15]. This has also been suspected by Liuba et al. in 2005 [9]. Their multiple hit theory states that repeated episodes of acute infections might enhance oxidized modification of LDL, which plays an important role in the development of atherosclerosis by fostering foam cell accumulation and subsequent thickening of the vascular wall [39, 40]. This association of inflammation and formation of atherogenic oxidized LDL (oxLDL) has been described in autoimmune disorders as well and a comparable pathomechanism is imaginable in allergic diseases [2, 3]. Acute allergic reactions cause IgE-triggered mast cell activation and an acute-phase reaction, both leading to increased oxidative stress [2, 18, 41]. Activated mast cells degranulate cytokines, leukotrienes, prostaglandins and histamine, leading to endothelial activation and facilitated intracellular penetration of LDL [41]. Endothelial oxidative modification of LDL might be enhanced in children with allergic diseases, due to increased oxidative stress and decreased antioxidative capacity [2, 42]. Furthermore, several authors suggested that IgE-triggered mast cell activation during acute allergic reactions might lead to facilitated presentation of LDL to macrophages subsequently enhancing formation of foam cells [43, 44]. It was also suggested, that atherogenic complexes of CRP and oxLDL with or without β 2-glycoprotein-I might be present during acute phase reactions [2, 3, 45, 46]. Accordingly, the severity of clinical penetrance of atopic sensitization in terms of cumulative lifetime load of number and severity of allergic bouts might be a key factor to produce a relevant atopy-associated effect on vascular ageing. The results of our analyses on participants with allergic diseases do not support this hypothesis. However, we did not have information about the severity of past clinical allergic episodes in our participants and the lifetime load of antiallergic treatment. Yet, clarification of this question might be of importance for clinicians, as they would be encouraged to control allergic diseases very carefully in order to avoid adverse effects on long-term vascular health in affected children and adolescents. Future studies should therefore obtain detailed information about the number and severity of acute clinical exacerbations of allergic diseases of their participants in the past and address this question. Finally, it would be intriguing to compare the levels of oxLDL during chronic and acute hyperinflammatory activity in atopic children and adolescents and investigate their association with accelerated arterial stiffening and increased intima-media thickness.

Strengths and limitations

The standardized, high-quality measurements of DC and cIMT in a cohort of adolescents was a main strength of our study. Although early structural changes due to chronic systemic

inflammation have been reported in children and adolescents [15, 20, 47], one should be aware, that the assessment of functional instead of structural alterations (e.g. by flow-mediated dilation), might lead to earlier detection of increased vascular risk related to atopic sensitization. Furthermore, our study does not allow establishing causality, due to its cross-sectional nature. Yet, there was a tendency towards an association of atopic sensitization with decreased DC and the non-significance of the results might be, at least in part, due to the relatively small sample size. We performed analysis of total and specific IgE, which are qualitative markers of the degree of atopic sensitization on the immunological level [38]. Further, based on interview and clinical measurements, our cohort had a relatively thorough phenotyping with regard to allergy. Further detail about the number and severity of clinical episodes of allergic disease, earlier medical treatment and assessment of further inflammatory markers (i. e. oxLDL, high-sensitivity C-reactive protein, soluble interleukin-2 receptor, eosinophil cationic protein) would have been helpful for a characterization of the lifetime load of hyperinflammatory activity in our study population. This may be especially relevant, as the relatively small number of participants with symptomatic allergic disease might be a major limiting factor for the non-significance of our results. Future studies should aim towards a detailed characterization of the extent and severity of inflammatory activity in their participants, as well as a larger sample size. A selection bias due to the exclusion of the 21 participants, due to denied parental consent for blood tests, seems also very unlikely, as their empirical, vascular and clinical allergy characteristics were not different from those included in the study.

Conclusions and perspectives

To the best of our knowledge, this study is the first to analyze a possible association of atopic sensitization and allergic diseases with DC and cIMT in adolescents. Whereas evidence points towards an impact of systemic hyper-inflammation due to atopic sensitization on the vascular endothelium, our results do not support this assumption in adolescents [2, 3, 9, 15, 20, 39]. Better knowledge about the impact of the clinical character, main determinants and potential role of disease control of either atopic sensitization or allergic diseases might be valuable. Further studies should investigate, whether the number and severity of repeated acute clinical bouts of allergic diseases might be a predictor of early vascular ageing, rather than chronic low-grade inflammatory activity mediated by atopic sensitization.

Supporting information

S1 File. Definition of “atopic sensitization” and “allergic disease”, anthropometric and physical activity measures.

(PDF)

S2 File. Calculation of “DC” and “cIMT”.

(PDF)

S1 Dataset.

(SAV)

Acknowledgments

We would like to wholeheartedly thank the study participants and support staff that facilitated the study. We would also thank Professor Andreas Bircher (Department of Dermatology, Allergy Unit, University Hospital Basel, Switzerland) for his valuable expert advice.

The authors declare that they have no conflicts of interest or sources of income relating to the research. The data are presented clearly, honestly, and without fabrication, falsification or inappropriate manipulation.

Author Contributions

Conceptualization: Ane Johannessen, Cecilie Svanes, Julia Dratva.

Data curation: Karsten Königstein, Denis Infanger, Randi Jacobsen Bertelsen.

Formal analysis: Karsten Königstein, Denis Infanger, Randi Jacobsen Bertelsen.

Funding acquisition: Ane Johannessen.

Investigation: Randi Jacobsen Bertelsen, Ane Johannessen.

Methodology: Randi Jacobsen Bertelsen, Ulrike Waje-Andreassen, Arno Schmidt-Trucksäss, Julia Dratva.

Project administration: Cecilie Svanes.

Resources: Ane Johannessen, Ulrike Waje-Andreassen, Julia Dratva.

Software: Denis Infanger, Ulrike Waje-Andreassen, Arno Schmidt-Trucksäss.

Supervision: Ulrike Waje-Andreassen, Arno Schmidt-Trucksäss, Cecilie Svanes, Julia Dratva.

Validation: Arno Schmidt-Trucksäss, Cecilie Svanes, Julia Dratva.

Visualization: Julia Dratva.

Writing – original draft: Karsten Königstein.

Writing – review & editing: Karsten Königstein, Randi Jacobsen Bertelsen, Ane Johannessen, Ulrike Waje-Andreassen, Arno Schmidt-Trucksäss, Cecilie Svanes, Julia Dratva.

References

1. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *The New England journal of medicine*. 1997; 336(14):973–9. <https://doi.org/10.1056/NEJM199704033361401> PMID: 9077376
2. Levitan I, Volkov S, Subbaiah PV. Oxidized LDL: diversity, patterns of recognition, and pathophysiology. *Antioxidants & redox signaling*. 2010; 13(1):39–75.
3. Matsuura E, Hughes GR, Khamashta MA. Oxidation of LDL and its clinical implication. *Autoimmun Rev*. 2008; 7(7):558–66. <https://doi.org/10.1016/j.autrev.2008.04.018> PMID: 18625445
4. Wang J, Cheng X, Xiang MX, Alanne-Kinnunen M, Wang JA, Chen H, et al. IgE stimulates human and mouse arterial cell apoptosis and cytokine expression and promotes atherogenesis in Apoe^{-/-} mice. *The Journal of clinical investigation*. 2011; 121(9):3564–77. <https://doi.org/10.1172/JCI46028> PMID: 21821913
5. Marciniak-Sroka J, Jawien J, Jakiela B, Szczeklik A. Increase in plasma sCD23 levels precedes immunoglobulin E elevation after coronary artery bypass graft surgery. *Polskie Archiwum Medycyny Wewnętrznej*. 2011; 121(4):109–14. PMID: 21532532
6. Kovanen PT. Mast cells: multipotent local effector cells in atherothrombosis. *Immunological reviews*. 2007; 217:105–22. <https://doi.org/10.1111/j.1600-065X.2007.00515.x> PMID: 17498055
7. Wick G, Knoflach M, Xu Q. Autoimmune and inflammatory mechanisms in atherosclerosis. *Annual review of immunology*. 2004; 22:361–403. <https://doi.org/10.1146/annurev.immunol.22.012703.104644> PMID: 15032582
8. Dwyer JH, Allayee H, Dwyer KM, Fan J, Wu H, Mar R, et al. Arachidonate 5-lipoxygenase promoter genotype, dietary arachidonic acid, and atherosclerosis. *The New England journal of medicine*. 2004; 350(1):29–37. <https://doi.org/10.1056/NEJMoa025079> PMID: 14702425
9. Liuba P, Pesonen E. Infection and early atherosclerosis: does the evidence support causation? *Acta Paediatr*. 2005; 94(6):643–51. <https://doi.org/10.1111/j.1651-2227.2005.tb01958.x> PMID: 16188762

10. Berenson GS. Childhood risk factors predict adult risk associated with subclinical cardiovascular disease. The Bogalusa Heart Study. *The American journal of cardiology*. 2002; 90(10C):3L–7L. [https://doi.org/10.1016/s0002-9149\(02\)02953-3](https://doi.org/10.1016/s0002-9149(02)02953-3) PMID: 12459418
11. Ferreira I, van de Laar RJ, Prins MH, Twisk JW, Stehouwer CD. Carotid stiffness in young adults: a life-course analysis of its early determinants: the Amsterdam Growth and Health Longitudinal Study. *Hypertension*. 2012; 59(1):54–61. <https://doi.org/10.1161/HYPERTENSIONAHA.110.156109> PMID: 22068867
12. Rosenfeld ME, Campbell LA. Pathogens and atherosclerosis: update on the potential contribution of multiple infectious organisms to the pathogenesis of atherosclerosis. *Thrombosis and haemostasis*. 2011; 106(5):858–67. <https://doi.org/10.1160/TH11-06-0392> PMID: 22012133
13. Celermajer DS, Adams MR, Clarkson P, Robinson J, McCredie R, Donald A, et al. Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. *The New England journal of medicine*. 1996; 334(3):150–4. <https://doi.org/10.1056/NEJM199601183340303> PMID: 8531969
14. Dalla Pozza R, Bechtold S, Bonfig W, Putzker S, Kozlik-Feldmann R, Netz H, et al. Age of onset of type 1 diabetes in children and carotid intima medial thickness. *The Journal of clinical endocrinology and metabolism*. 2007; 92(6):2053–7. <https://doi.org/10.1210/jc.2006-2868> PMID: 17374703
15. Dratva J, Caviezel S, Schaffner E, Bettschart R, Kuenzli N, Schindler C, et al. Infectious diseases are associated with carotid intima media thickness in adolescence. *Atherosclerosis*. 2015; 243(2):609–15. <https://doi.org/10.1016/j.atherosclerosis.2015.10.021> PMID: 26545015
16. Elkiran O, Yilmaz E, Koc M, Kamanli A, Ustundag B, Ilhan N. The association between intima media thickness, central obesity and diastolic blood pressure in obese and overweight children: a cross-sectional school-based study. *International journal of cardiology*. 2013; 165(3):528–32. <https://doi.org/10.1016/j.ijcard.2011.09.080> PMID: 22014414
17. Knoflach M, Kiechl S, Mayr A, Willeit J, Poewe W, Wick G. Allergic rhinitis, asthma, and atherosclerosis in the Bruneck and ARMY studies. *Archives of internal medicine*. 2005; 165(21):2521–6. <https://doi.org/10.1001/archinte.165.21.2521> PMID: 16314550
18. Potaczek DP. Links between allergy and cardiovascular or hemostatic system. *International journal of cardiology*. 2014; 170(3):278–85. <https://doi.org/10.1016/j.ijcard.2013.11.029> PMID: 24315352
19. Cakmak A, Zeyrek D, Cece H, Erel O. The relationship between carotid intima media thickness and oxidative stress in asthmatic children. *Asian Pac J Allergy Immunol*. 2010; 28(4):256–61. PMID: 21337909
20. Evelein AM, Visseren FL, van der Ent CK, Grobbee DE, Uiterwaal CS. Allergies are associated with arterial changes in young children. *European journal of preventive cardiology*. 2015; 22(11):1480–7. <https://doi.org/10.1177/2047487314554863> PMID: 25301873
21. Teynor A, Caviezel S, Dratva J, Kunzli N, Schmidt-Trucksass A. An automated, interactive analysis system for ultrasound sequences of the common carotid artery. *Ultrasound in medicine & biology*. 2012; 38(8):1440–50.
22. Touboul PJ, Hennerici MG, Meairs S, Adams H, Amarenco P, Bornstein N, et al. Mannheim carotid intima-media thickness and plaque consensus (2004–2006–2011). An update on behalf of the advisory board of the 3rd, 4th and 5th watching the risk symposia, at the 13th, 15th and 20th European Stroke Conferences, Mannheim, Germany, 2004, Brussels, Belgium, 2006, and Hamburg, Germany, 2011. *Cerebrovascular diseases*. 2012;34(4):290–6.
23. Touboul PJ, Prati P, Scarabin PY, Adrai V, Thibout E, Ducimetiere P. Use of monitoring software to improve the measurement of carotid wall thickness by B-mode imaging. *Journal of hypertension Supplement: official journal of the International Society of Hypertension*. 1992; 10(5):S37–41.
24. Dalla Pozza R, Ehringer-Schetitska D, Fritsch P, Jokinen E, Petropoulos A, Oberhoffer R, et al. Intima media thickness measurement in children: A statement from the Association for European Paediatric Cardiology (AEPIC) Working Group on Cardiovascular Prevention endorsed by the Association for European Paediatric Cardiology. *Atherosclerosis*. 2015; 238(2):380–7. <https://doi.org/10.1016/j.atherosclerosis.2014.12.029> PMID: 25555270
25. Doyon A, Kracht D, Bayazit AK, Deveci M, Duzova A, Krmar RT, et al. Carotid artery intima-media thickness and distensibility in children and adolescents: reference values and role of body dimensions. *Hypertension*. 2013; 62(3):550–6. <https://doi.org/10.1161/HYPERTENSIONAHA.113.01297> PMID: 23817494
26. Endes S, Caviezel S, Schaffner E, Dratva J, Schindler C, Kunzli N, et al. Associations of Novel and Traditional Vascular Biomarkers of Arterial Stiffness: Results of the SAPALDIA 3 Cohort Study. *PloS one*. 2016; 11(9):e0163844. <https://doi.org/10.1371/journal.pone.0163844> PMID: 27685325
27. Lamotte C, Iliescu C, Libersa C, Gottrand F. Increased intima-media thickness of the carotid artery in childhood: a systematic review of observational studies. *Eur J Pediatr*. 2011; 170(6):719–29. <https://doi.org/10.1007/s00431-010-1328-y> PMID: 20978785

28. van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R. *Journal of statistical software*. 2011; 45(3):1–67.
29. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Statistics in medicine*. 2011; 30(4):377–99. <https://doi.org/10.1002/sim.4067> PMID: 21225900
30. van Buuren S. *Flexible imputation of missing data*. Boca Raton, FL: CRC Press; 2012.
31. Barnard J, Rubin DB. Small-sample degrees of freedom with multiple imputation. *Biometrika*. 1999; 86(4):948–55.
32. Kallio K, Jokinen E, Saarinen M, Hamalainen M, Volanen I, Kaitosaari T, et al. Arterial intima-media thickness, endothelial function, and apolipoproteins in adolescents frequently exposed to tobacco smoke. *Circ Cardiovasc Qual Outcomes*. 2010; 3(2):196–203. <https://doi.org/10.1161/CIRCOUTCOMES.109.857771> PMID: 20197510
33. Goksor E, Amark M, Alm B, Gustafsson PM, Wennergren G. The impact of pre- and post-natal smoke exposure on future asthma and bronchial hyper-responsiveness. *Acta Paediatr*. 2007; 96(7):1030–5. <https://doi.org/10.1111/j.1651-2227.2007.00296.x> PMID: 17498194
34. Rasmussen F, Lambrechtsen J, Siersted HC, Hansen HS, Hansen NC. Low physical fitness in childhood is associated with the development of asthma in young adulthood: the Odense schoolchild study. *Eur Respir J*. 2000; 16(5):866–70. PMID: 11153585
35. Skilton MR, Viikari JS, Juonala M, Laitinen T, Lehtimäki T, Taittonen L, et al. Fetal growth and preterm birth influence cardiovascular risk factors and arterial health in young adults: the Cardiovascular Risk in Young Finns Study. *Arterioscler Thromb Vasc Biol*. 2011; 31(12):2975–81. <https://doi.org/10.1161/ATVBAHA.111.234757> PMID: 21940950
36. Horta BL, Gigante DP, Lima RC, Barros FC, Victora CG. Birth by caesarean section and prevalence of risk factors for non-communicable diseases in young adults: a birth cohort study. *PloS one*. 2013; 8(9): e74301. <https://doi.org/10.1371/journal.pone.0074301> PMID: 24040224
37. Juliusson PB, Roelants M, Nordal E, Furevik L, Eide GE, Moster D, et al. Growth references for 0–19 year-old Norwegian children for length/height, weight, body mass index and head circumference. *Annals of human biology*. 2013; 40(3):220–7. <https://doi.org/10.3109/03014460.2012.759276> PMID: 23414181
38. Sicherer SH, Wood RA, American Academy of Pediatrics Section On A, Immunology. Allergy testing in childhood: using allergen-specific IgE tests. *Pediatrics*. 2012; 129(1):193–7. <https://doi.org/10.1542/peds.2011-2382> PMID: 22201146
39. Liuba P, Persson J, Luoma J, Yla-Herttuala S, Pesonen E. Acute infections in children are accompanied by oxidative modification of LDL and decrease of HDL cholesterol, and are followed by thickening of carotid intima-media. *European heart journal*. 2003; 24(6):515–21. [https://doi.org/10.1016/s0195-668x\(02\)00750-9](https://doi.org/10.1016/s0195-668x(02)00750-9) PMID: 12643884
40. Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *The New England journal of medicine*. 1989; 320(14):915–24. <https://doi.org/10.1056/NEJM198904063201407> PMID: 2648148
41. Togias A. Systemic effects of local allergic disease. *J Allergy Clin Immunol*. 2004; 113(1 Suppl):S8–14.
42. Zeyrek D, Cakmak A, Atas A, Kocyigit A, Erel O. DNA damage in children with asthma bronchiale and its association with oxidative and antioxidative measurements. *Pediatr Allergy Immunol*. 2009; 20(4):370–6. <https://doi.org/10.1111/j.1399-3038.2008.00780.x> PMID: 18801009
43. Ma H, Kovanen PT. IgE-dependent generation of foam cells: an immune mechanism involving degranulation of sensitized mast cells with resultant uptake of LDL by macrophages. *Arterioscler Thromb Vasc Biol*. 1995; 15(6):811–9. PMID: 7773738
44. Ma H, Kovanen PT. Inhibition of mast cell-dependent conversion of cultured macrophages into foam cells with antiallergic drugs. *Arterioscler Thromb Vasc Biol*. 2000; 20(12):E134–42. PMID: 11116078
45. Chang MK, Binder CJ, Torzewski M, Witztum JL. C-reactive protein binds to both oxidized LDL and apoptotic cells through recognition of a common ligand: Phosphorylcholine of oxidized phospholipids. *Proceedings of the National Academy of Sciences of the United States of America*. 2002; 99(20):13043–8. <https://doi.org/10.1073/pnas.192399699> PMID: 12244213
46. Tabuchi M, Inoue K, Usui-Kataoka H, Kobayashi K, Teramoto M, Takasugi K, et al. The association of C-reactive protein with an oxidative metabolite of LDL and its implication in atherosclerosis. *J Lipid Res*. 2007; 48(4):768–81. <https://doi.org/10.1194/jlr.M600414-JLR200> PMID: 17261875
47. Fernberg U, Fernstrom M, Hurtig-Wennlof A. Arterial stiffness is associated to cardiorespiratory fitness and body mass index in young Swedish adults: The Lifestyle, Biomarkers, and Atherosclerosis study. *European journal of preventive cardiology*. 2017; 24(17):1809–18. <https://doi.org/10.1177/2047487317720796> PMID: 28696134